

学校编码: 10384

学号: 22320130153789

分类号__密级__

UDC__

厦门大学

博 士 学 位 论 文

珊瑚共生体对 UV 辐射与 CO₂ 浓度变化
的生理学响应

Physiological Responses of Coral Symbiont to Changes of
UV radiation and CO₂ concentrations

周洁

指导教师姓名: 高坤山 教授

专 业 名 称: 海洋生物学

论文提交日期: 2016 年 11 月

论文答辩时间: 2016 年 11 月

学位授予日期: 2016 年

答辩委员会主席: __

评阅人: __

2016 年 11 月

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下,独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果,均在文中以适当方式明确标明,并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外,该学位论文为(高坤山 教授)课题(组)的研究成果,获得(高坤山 教授)课题(组)经费或实验室的资助,在(海洋环境变化生理学)实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称,未有此项声明内容的,可以不作特别声明。)

声明人(签名):

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文（包括纸质版和电子版），允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于：

（ ） 1. 经厦门大学保密委员会审查核定的保密学位论文，
于 年 月 日解密，解密后适用上述授权。

（ ☒ ） 2. 不保密，适用上述授权。

（请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文，未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的，默认为公开学位论文，均适用上述授权。）

声明人（签名）：

年 月 日

目录

摘要.....	I
Abstract.....	IV
缩略词 (Abbreviations)	VII
第一章 文献综述与研究意义	1
1.1 珊瑚及其共生藻的生物学研究进展	1
1.1.1 共生藻 (<i>Symbiodinium</i> sp.) 的生物学特性.....	1
1.1.2 珊瑚与共生藻间共生关系的建立	3
1.1.3 珊瑚与共生藻间的共生	4
1.1.4 珊瑚共生体的碳浓缩机制	6
1.2 环境因子对珊瑚共生体的影响	8
1.2.1 温度对珊瑚共生体的影响	9
1.2.2 紫外辐射对珊瑚共生体的影响	10
1.2.3 海洋酸化对珊瑚共生体的影响	12
1.3 研究方法	17
1.3.1 紫外辐射效应的研究方法	17
1.3.2 海洋酸化效应的研究方法	18
1.3.2.1 自然海区的监测	20
1.3.2.2 控制培养体系	21
1.4 研究目的与意义	22
第二章 紫外辐射对珊瑚共生藻日释放量的影响	24
2.1 前言	24
2.2 材料与方法	25
2.2.1 珊瑚采集	25
2.2.2 紫外辐射处理	25
2.2.3 实验设计	26
2.2.4 光化学效率的测定	27
2.2.5 共生藻释放速率的测定	27
2.2.6 珊瑚样品中共生藻密度与色素浓度的测定	27
2.2.7 色素含量与紫外吸收物质 (UVACs) 的测定	28
2.2.8 数据分析	29
2.3 结果	29
2.3.1 共生藻光化学效率的日变化	29
2.3.2 珊瑚共生藻的日间释放模式	31
2.3.3 色素的日间释放模式	31

2.3.4 珊瑚共生藻在日间的总释放量	31
2.3.5 珊瑚样品中色素含量与共生藻密度	34
2.4 讨论	36
第三章 紫外辐射对珊瑚幼体及其共生藻的影响	38
第一节 紫外辐射对鹿角杯形珊瑚 (<i>Pocillopora damicornis</i>) 幼体及其共生藻的影响	38
3.1.1 前言	38
3.1.2 材料与方法	39
3.1.2.1 珊瑚的采集及幼体收集	39
3.1.2.2 实验设计	40
3.1.2.3 珊瑚幼体成活率、附着率和变态率的测定	43
3.1.2.4 珊瑚幼体中色素含量与紫外吸收物质 (UVACs) 的测定	43
3.1.2.5 珊瑚幼体中共生藻密度的测定	44
3.1.2.6 光合放氧和暗呼吸的测定	44
3.1.2.7 光化学效率的测定	45
3.1.2.8 数据分析	45
3.1.3 结果	45
3.1.3.1 鹿角杯形珊瑚 (<i>Pocillopora damicornis</i>) 幼体的生长发育	45
3.1.3.2 鹿角杯形珊瑚 (<i>Pocillopora damicornis</i>) 幼体色素含量与共生藻密度	47
3.1.3.3 光合放氧与暗呼吸速率	51
3.1.4 讨论	53
第二节 紫外辐射对浅杯排孔珊瑚 (<i>Seriatopora caliendrum</i>) 幼体及其共生藻的影响	56
3.2.1 前言	56
3.2.2 材料与方法	56
3.2.2.1 珊瑚的采集及幼体收集	57
3.2.2.2 实验设计	57
3.2.2.3 珊瑚幼体成活率、附着率和变态率的测定	59
3.2.2.4 珊瑚幼体中色素含量与紫外吸收物质 (UVACs) 的测定	59
3.2.2.5 珊瑚幼体中共生藻密度的测定	59
3.2.2.6 光合放氧和暗呼吸的测定	59
3.2.2.7 光化学效率的测定	59
3.2.2.8 数据分析	59
3.2.3 结果	60
3.2.3.1 浅杯排孔珊瑚 (<i>Seriatopora caliendrum</i>) 幼体的生长发育	60
3.2.3.2 浅杯排孔珊瑚 (<i>Seriatopora caliendrum</i>) 幼体色素含量与共生藻密度	61
3.2.3.3 光合放氧与暗呼吸速率	63
3.2.4 讨论	65

第四章 共生藻对 CO₂ 昼降夜升的生理学响应68

4.1 前言.....68

4.2 材料与方法70

4.2.1 藻种及培养条件70

4.2.2 海水碳酸盐系统的调控71

4.2.3 比生长速率和细胞色素含量的测定71

4.2.4 叶绿素荧光参数的测定72

4.2.5 无机碳亲和力的测定73

4.2.6 光合放氧和暗呼吸的测定74

4.2.7 胞内碳库的测定74

4.2.8 数据分析75

4.3 结果.....76

4.3.1 碳酸盐系统76

4.3.2 对生长和细胞色素含量的影响77

4.3.3 对光化学特性的影响78

4.3.4 对光合放氧和呼吸速率的影响80

4.3.5 对无机碳亲和力和胞内碳库的影响81

4.4 讨论.....84

第五章 共生藻对 CO₂ 浓度升高与紫外辐射耦合效应的生理学响应87

5.1 前言.....87

5.2 材料与方法88

5.2.1 藻种及培养条件88

5.2.2 海水碳酸盐系统的调控88

5.2.3 紫外辐射处理89

5.2.4 比生长速率和细胞色素含量的测定89

5.2.5 叶绿素荧光参数的测定89

5.2.6 无机碳亲和力的测定89

5.2.7 呼吸速率的测定89

5.2.8 数据分析89

5.3 结果.....90

5.3.1 碳酸盐系统90

5.3.2 不同 $p\text{CO}_2$ 处理下共生藻光化学效率对短期紫外辐射的响应91

5.3.3 对生长和细胞色素含量的影响95

5.3.4 对光化学特性的影响98

5.3.5 对无机碳亲和力的影响101

5.3.6 对呼吸速率的影响103

5.4 讨论.....104

总结与创新点	108
参考文献.....	111
论文发表情况	144
致谢.....	145

厦门大学博硕士论文摘要库

Table of Contents

Abstract (In Chinese).....	I
Abstract (In English)	IV
Abbreviations	VII
Chapter 1 Introduction.....	1
1.1 Research progress on the biology of Coral-Dinoflagellate symbiosis.....	1
1.1.1 Biology of <i>Symbiodinium</i> sp.	1
1.1.2 Establishment of coral-algal symbiosis	3
1.1.3 Cost-benefit analysis of the symbiosis	4
1.1.4 Carbon concentrating mechanism (CCMs)	6
1.2 Environmental factors on Coral-Dinoflagellate symbiosis	8
1.2.1 Temperature	9
1.2.2 Ultraviolet radiation	10
1.2.3 Ocean acidification	12
1.3 Methodology	17
1.3.1 Methods for studying UVR effects.....	17
1.3.2 Methods for ocean acidification researches	18
1.4 Research objectives and scheme	22
Chapter 2 Effect of UV radiation on <i>Symbiodinium</i> release in coral <i>Pocillopora damicornis</i>	24
2.1 Introduction.....	24
2.2 Materials and Methods.....	25
2.2.1 Sampling of coral colony	25
2.2.2 UV radiation treatments	25
2.2.3 Experimental setup	26
2.2.4 Chlorophyll fluorescence measurements.....	27
2.2.5 Measurements of <i>Symbiodinium</i> release rate	27
2.2.6 Determination of <i>Symbiodinium</i> density and pigment content.....	27
2.2.7 Determination of pigment and UVACs content.....	28
2.2.8 Statistical analysis	29
2.3 Results	29
2.3.1 Diurnal photosynthetic efficiency of <i>Symbiodinium</i>	30
2.3.2 Day-time expulsion pattern of <i>Symbiodinium</i>	31
2.3.3 Day-time expulsion pattern of pigments	31
2.3.4 Total release amount of <i>Symbiodinium</i>	31
2.3.5 Photosynthetic pigments and <i>Symbiodinium</i> densities in coral nubbins ..	34
2.4 Discussion.....	36
Chapter 3 Incident ultraviolet irradiances influence on the coral	

larva and its symbiotic algae.....38

3.1 Effects of UV on the larval symbiont of coral <i>Pocillopora damicornis</i>	38
3.1.1 Introduction	38
3.1.2 Materials and Methods	39
3.1.2.1 Colony sampling and coral larvae collection.....	39
3.1.2.2 Experimental setup.....	40
3.1.2.3 Mortality and development measurements	43
3.1.2.4 Determination of pigment and UVACs content.....	43
3.1.2.5 Determination of <i>Symbiodinium</i> density.....	44
3.1.2.6 Determination of photosynthetic oxygen evolution and respiration	44
3.1.2.7 Chlorophyll fluorescence measurements	45
3.1.2.8 Statistical analysis	45
3.1.3 Results	45
3.1.3.1 Development of coral larvae	45
3.1.3.2 Pigment content and <i>Symbiodinium</i> density	47
3.1.3.3 Photosynthetic rate and respiration	51
3.1.4 Discussion	53
3.2 Effects of UV on the larval symbiont of coral <i>Seriatopora caliendrum</i>	56
3.2.1 Introduction	56
3.2.2 Materials and Methods	56
3.2.2.1 Colony sampling and coral larvae collection.....	57
3.2.2.2 Experimental setup.....	57
3.2.2.3 Mortality and development measurements	59
3.2.2.4 Determination of pigment and UVACs content.....	59
3.2.2.5 Determination of <i>Symbiodinium</i> density.....	59
3.2.2.6 Determination of photosynthetic oxygen evolution and respiration	59
3.2.2.7 Chlorophyll fluorescence measurements	59
3.2.2.8 Statistical analysis	59
3.2.3 Results	60
3.2.3.1 Development of coral larvae	60
3.2.3.2 Pigment content and <i>Symbiodinium</i> density	61
3.2.3.3 Photosynthetic rate and respiration	63
3.2.4 Discussion	65

Chapter 4 Physiological responses of *Symbiodinium* to diurnal oscillating pCO₂.....68

4.1 Introduction.....	68
4.2 Materials and Methods.....	70
4.2.1 Species and culture conditions	70
4.2.2 Perturbation of carbonate system	71
4.2.3 Determination of growth and pigments	71
4.2.4 Chlorophyll fluorescence measurements.....	72
4.2.5 Determination of photosynthetic inorganic carbon affinity	73
4.2.6 Determination of photosynthetic oxygen evolution and respiration	74
4.2.7 Determination of intracellular carbon concentration.....	74
4.2.8 Statistical analysis	75
4.3 Results	76

4.3.1 Carbonate chemistry system	76
4.3.2 Growth and pigments	77
4.3.3 Chlorophyll fluorescence parameters	78
4.3.4 Photosynthetic rate and respiration	80
4.3.5 Photosynthetic inorganic carbon affinity and intracellular carbon concentration	81
4.4 Discussion.....	84
 Chapter 5 Combined effects of ocean acidification and UVR	87
5.1 Introduction.....	87
5.2 Materials and Methods.....	88
5.2.1 Species and culture conditions	88
5.2.2 Perturbation of carbonate system	88
5.2.3 UV radiation treatments	89
5.2.4 Determination of growth and pigments	89
5.2.5 Chlorophyll fluorescence measurements.....	89
5.2.6 Determination of photosynthetic inorganic carbon affinity	89
5.2.7 Determination of respiration	89
5.2.8 Statistical analysis	89
5.3 Results	90
5.3.1 Carbonate chemistry system.....	90
5.3.2 Chlorophyll fluorescence parameters during UVR exposure.....	91
5.3.3 Growth and pigments	95
5.3.4 Chlorophyll fluorescence parameters upon UVR exposure	98
5.3.5 Photosynthetic inorganic carbon affinity.....	101
5.3.6 Dark respiration.....	103
5.4 Discussion.....	104
 Conclusion and innovation.....	108
References	111
Publications	144
Acknowledgements	145

厦门大学博硕士论文摘要库

摘要

大气 CO₂ 浓度升高导致了全球变暖与海洋酸化, 并引起海水中有色可溶性有机物减少, 使得海水透明度增加, 提高了进入水体的紫外辐射 (UVR, 280—400 nm) 水平, 从而对浅水珊瑚礁生态系统的各种生物过程造成影响。另一方面, 海洋酸化及环境中 pH/CO₂ 的扰动, 也会影响珊瑚及其与藻类的共生关系。本博士论文, 以鹿角杯形珊瑚 (*Pocillopora damicornis*) 为研究对象, 探讨了 UV 辐射对其共生藻日释放量的影响; 同时, 研究了鹿角杯形珊瑚和浅杯排孔珊瑚 (*Seriatopora caliendrum*) 浮浪幼体对阳光 UV 辐射变化的响应。在此基础上, 揭示了 UV 辐射对浮浪幼体及其共生藻的影响。另外, 还研究了一种系群 D 共生藻 (CCMP2556) 对海水中 *p*CO₂ 昼降夜升的生理学响应, 并探讨了 *p*CO₂ 升高与 UVR 对其的复合效应。主要的结果如下:

1. 阳光辐射变化影响鹿角杯形珊瑚释放共生藻的量。UV 辐射显著降低了鹿角杯形珊瑚色素 (包括叶绿素 *a* 和类胡萝卜素) 和紫外吸收物质 (UVACs) 的含量, 导致珊瑚共生藻的日平均释放速率提高了 10—20% (95% 置信区间上不显著)。有 UV 和无 UV 辐射处理间的日变化模式相似, 即正午时共生藻的释放达到最高峰, 但 UV-B (280—315 nm) 的存在使得这一高峰期提前了 1 小时。
2. 不同种类的珊瑚幼体对 UV 辐射的响应不同。与分布较深的浅杯排孔珊瑚相比, 分布较浅的鹿角杯形珊瑚的浮浪幼体, 对 UV 辐射的敏感性高, 幼体内共生藻光合作用受抑制的程度也较大。在 UV 辐射的作用下, 鹿角杯形珊瑚幼体共生藻的净光合速率下降了 75—85%, 其叶绿素 *a* 和 UVAC 含量与温度变化的相关性大。UV-A (315—400 nm) 显著抑制了鹿角杯形珊瑚幼体的发育, 其对幼体的成活率、变态率与附着率的抑制率达 22—25%。浅杯排孔珊瑚幼体内的共生藻平均密度, 受 UV 辐射的影响, 反而升高。UV-A 辐射促进了共生藻的有效光化学效率, 而 UV-B 可抵消 UV-A 的促进作用; UV-A 降低了该珊瑚幼体的变态速率。
3. 海水 *p*CO₂ 升高和昼降夜升 (460—1000 μ atm) 都使得共生藻对无机碳的浓缩能力 (CCMs) 发生了上调, 但二者间有着不同的上调模式。*p*CO₂ 的增加促

进了共生藻细胞的比生长速率,降低了细胞内叶绿素 *a* 和类胡萝卜素的含量。同时,共生藻 PSII 的有效光化学截面积减小了,且 Q_A 的周转时间延长了,表观光能利用效率降低了;然而,其单位叶绿素 *a* 的光合放氧速率和呼吸速率分别提高了 49%和 41%;单位细胞的光合与呼吸速率均未受到影响。此外,*p*CO₂ 的增加使得 K_m(DIC)的值降低了 63%,而 K_m(CO₂)稳定在 3.0–3.4 μmol L⁻¹,提高了细胞对 HCO₃⁻的亲合力。*p*CO₂ 的昼降夜升降低了细胞内类胡萝卜素的含量。与 *p*CO₂ 的增加类似,*p*CO₂ 的昼降夜升使得 PSII 的有效光化学截面积减小,并延长了 Q_A 的周转时间,但 PSII 的有效光化学效率提高了 27%,非光化学淬灭下降了 34%,虽然如此,其光合放氧速率并未提高,且呼吸速率要明显低于高 CO₂ 处理 (19%)。此外,*p*CO₂ 的昼降夜升使得胞内碳库增大了近一倍。

4. *p*CO₂ 升高与 UV 辐射对共生藻有拮抗作用。在 UV 辐射处理下,共生藻细胞的捕光色素含量下降,而 *p*CO₂ 的增加使得其含量明显增加。同时,UV 辐射促使共生藻细胞 K_m(DIC)增加了 43–62%,K_m(CO₂)也相应提高了 43–64%,降低了细胞对无机碳的亲合力,而 *p*CO₂ 的增加造成细胞 K_m(DIC) 和 K_m(CO₂)都下降了 27–50%,即 CCMs 显著上调。同时,细胞的呼吸速率在二者的拮抗作用下,分别下降了 22–39%(LC: 低 CO₂ 处理)与增加了 33–36%(HC: 高 CO₂ 处理)。此外,在 UV 辐射下,HC 细胞对 UV-A 损伤的修复速率要明显高于 LC 细胞,但在培养光强下,LC 细胞对 UV-A 损伤的恢复速率要比 HC 细胞快了 33%。UV-A 对细胞的表观光能效率、最大电子传递速率和光饱和参数具有抑制作用,而 UV-B 的存在抵消了这一作用,*p*CO₂ 的增加使得细胞对 UV 的敏感性降低,缓解了紫外辐射的效应。

总之,UV 辐射,虽然对珊瑚共生藻的释放没有显著影响,但会抑制幼体的变态与附着过程,并抑制共生藻的光合作用。*p*CO₂ 的增加与昼降夜升都会提高共生藻对无机碳的利用能力,但在昼降夜升下光合或生长速率并未增加,说明 CO₂ 的扰动导致共生藻能耗增加。UV 辐射和 *p*CO₂ 升高产生了拮抗效应,高 CO₂ 下,细胞对 UV 的敏感性下降。然而,原位复杂环境下,分布不同海域、不同深度的珊瑚及共生藻,经受阳光 UV 辐射的程度不同,对 CO₂ 升高的响应,会存在较大的区域性差别。

关键词：珊瑚幼体；共生藻；紫外辐射；海洋酸化

厦门大学博士论文摘要库

Abstract

Rising atmospheric carbon dioxide (CO₂) is causing ocean acidification and ocean warming, with reduced amount of colored dissolved organic matter in the water. Ultimately, the coral reef water may become clearer and the transmission of ultraviolet radiation (UVR, 280-400 nm) into the oceans would be enhanced, which would affect various biological processes in shallow coral reef ecosystem. On the other hand, ocean acidification and the fluctuation in pH/pCO₂ may also influence the symbiosis between coral and the dinoflagellate *Symbiodinium*. In this study, the release rate of *Symbiodinium* from the coral *Pocillopora damicornis* was studied under UVR stress. Moreover, the responses of the planula larvae of *P. damicornis*, as well as that of *Seriatopora caliendrum*, to UV radiation were investigated, and the effects of UVR on the symbiotic algae within the larvae were examined. In addition, by adjusting pCO₂ fluctuating regimes (constant versus oscillatory), the effects of high pCO₂ and diel variation in pCO₂ on the physiological performance of *Symbiodinium* CCMP2556 were studied, and the combined effects of elevated CO₂ and UVR were also examined. The main results are as follows:

1. Solar radiation modulated the diurnal release pattern of *Symbiodinium* from the coral *P. damicornis*. The UVR exposure significantly reduced the content of chl *a* and carotenoids per surface area in the coral branch, as well as that of the UV-absorbing compound (UVAC) at the end of experiment. The release rate was higher in the exposure to UVR by ~10-20%, but this increment was not significant at the 95% confidence level. The diurnal release pattern of *Symbiodinium* with or without UVR were similar, i.e. the release rate peaked at noon (11:00-13:00), while the time was advanced for an hour when exposed to UV-A (315-400 nm) and UV-B (280-315 nm).
2. The larvae of two coral species responded differently to UVR stress, with larvae of *P. damicornis* from shallower water being more sensitive to UVR exposure than *S. caliendrum* from deeper water, so as the inhibition to the photosynthesis of *Symbiodinium* in the larvae. UVR stress decreased net photosynthetic rate of endosymbiotic alga in *P. damicornis* larvae by 75-85%. There was a positive relationship between significance of differences in chl *a* and temperature, which

also occurred between the concentration of UV absorbing compounds and temperature. The development of *P. damicornis* larvae was severely inhibited by UV-A, with the inhibition rate of survivorship, metamorphosis and settlement ranging around 22-25%. The density of *Symbiodinium* in *S. caliendrum* larvae exposed to UVR was obviously higher than the control. The effective photochemical efficiency (F_v'/F_m') of *Symbiodinium* in *S. caliendrum* larvae was stimulated by UV-A, but the stimulation was offset by UV-B. UV-A obviously delayed the metamorphosis and settlement of *S. caliendrum* larvae.

3. Increased $p\text{CO}_2$ and day-night fluctuation of $p\text{CO}_2$ up-regulated the carbon concentrating mechanisms (CCMs) of *Symbiodinium* (CCMP2556). After acclimation to high $p\text{CO}_2$ environment for over 15 generations, the *Symbiodinium* showed significant higher growth rate, while the content of chl *a* and carotenoids per cell decreased. At the same time, the functional absorption cross-section of PSII (Sigma(II)_{440}) and the apparent photosynthetic efficiency (α) derived from light curve in 440 nm decreased, with the increase of time constant of Q_A turnover (τ). The photosynthetic oxygen evolution and respiration rates per chl *a* were enhanced by 49% and 41%, respectively, whereas both rates normalized to per cell were unchanged. In view of the carbon utilization, increasing $p\text{CO}_2$ significantly decreased the half-saturation concentration for DIC ($K_m(\text{DIC})$) by 63%, while the $K_m(\text{CO}_2)$ stabilized at 3.0 to 3.4 $\mu\text{mol L}^{-1}$, and the affinity to HCO_3^- was thus enhanced. Variable $p\text{CO}_2$ (oscillating between 460/1000 μatm) caused the evident decrease of cellular carotenoids content, being similar with the effects of increasing $p\text{CO}_2$, Sigma(II)_{440} decreased and τ increased, while the effective quantum yield of PSII (F_v'/F_m') increased by 27% and non-photochemical quenching (NPQ) lowered by 34%. Moreover, the respiration rate of *Symbiodinium* was lower by 19% under diurnal fluctuating $p\text{CO}_2$, compared to that of the algae under increasing $p\text{CO}_2$ treatment. In face of variation of $p\text{CO}_2$, *Symbiodinium* up-regulated the CCMs by nearly doubling the internal carbon pool, without altering the carbon affinity.
4. An antagonistic effect occurred when elevated CO_2 was combined with UVR. Exposure to UVR reduced the cellular contents of chl *a* and chl *c*, but the inhibition was reversed when combined with increased $p\text{CO}_2$. In addition, upon UVR exposure, $K_m(\text{DIC})$ and $K_m(\text{CO}_2)$ of *Symbiodinium* increased by 43-62% and 43-

Degree papers are in the “[Xiamen University Electronic Theses and Dissertations Database](#)”.

Fulltexts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.